

Review Article	Pak-Euro Journal of Medical and Life Sciences
DOI: 10.31580/pjmls.v3i3.1621	Copyright © All rights are reserved by Corresponding Author
Vol. 3 No. 3, 2020: pp. 131-137	
www.readersinsight.net/pjmls	
Submission: September 03, 2020	Revised: September 26, 2020 Accepted: September 30, 2020

CAMPYLOBACTER-REVIEW ON ITS SIGNIFICANCE AS A FOODBORNE PATHOGEN

Tauseef Muhammad Asmat^{1*}, Aitbar Khan²

¹Center for Advanced Studies in Vaccinology and Biotechnology (CASVAB),

University of Balochistan, Quetta, Pakistan

²Balochistan Livestock & Dairy Development Department, Quetta, Pakistan

Corresponding Author: Tauseef M Asmat. E-mail: tauseefcasvab@gmail.com



Abstract

Food borne pathogens significantly cause illness in human which vary from mild symptoms to life threatening disease depending on the nature of microbial pathogens. Bacteria are almost responsible for 75% of food borne diseases in human. The food products especially of animal origin or their by products serve as the major source of transmission of food borne pathogens. All types of meat mainly beef, lamb and chicken and their by products can be contaminated with food borne pathogens like *Campylobacter*, *Salmonella*, *Staphylococcus*, *E. coli* and *Listeria*. These bacterial pathogens are equipped with several virulence factors and toxins which lead to illness in human. The severity of the disease mainly depends on the virulence of ingested pathogen. *Campylobacter* has been reported as the most frequent causative agent regarding foodborne illness. This article is aimed to review all aspects of *Campylobacter* as a foodborne pathogen.

Keywords: *Campylobacter*, *Campylobacteriosis*, Foodborne pathogen, Zoonotic agent, Food Contamination

INTRODUCTION

Although the food storage and its consumption has been greatly modified in recent decades and strict hygienic measures are followed to prevent any sort of contamination, still food borne pathogens pose a great health risk worldwide especially in developing countries (1). Almost, 600 million people each year suffer with food borne sickness around the globe and almost 2 million deaths are estimated only in developing countries because of this illness (2). This overwhelming sickness and deaths have great impact on health system and economy (3). The major foods borne out breaks are caused by different pathogens including bacteria, viruses and parasites (4). The food can be contaminated by these pathogens during slaughtering, harvesting, cooking and storage. In most of the cases the origin of these pathogens is from animals and can be transmitted to human when processed for food consumption (5). The other sources of food contamination are handlers, contaminated water, working place/utensils and in some case the air and dust (6).

The developed countries have significantly minimized the risk of outbreaks caused by food borne pathogens by implementing strict hygienic practices, modified the food safety and security laws and proper certification of the food handlers and restaurants (7-

11). However, such practices have not been adopted in developing countries and the food quality in terms of free from pathogens is not that promising (12-17). Strict implementation of food safety laws and public awareness are the only solutions to minimize the risk of food contamination with microbial pathogens.

This review article summarizes the importance, prevalence, disease outbreak and pathogenesis of *Campylobacter*.

CAMPYLOBACTER

Infection caused by *Campylobacter* is known as campylobacteriosis (18). This is known as the most common cause of diarrhea caused by any bacteria (19). *Campylobacter* are classified as Gram-negative and are curved or rod shaped bacteria (20). These are equipped with either single or double polar flagella and in some species they lack the flagellum. *Campylobacter* are micro-aerobic and needs special conditions to grow and don't produce spores (20). Campylobacteriosis, the disease caused by *Campylobacter* has been reported worldwide and has increased significantly during recent decades (21). Only in USA 20 out of 100,000 people suffer with this disease, however significant number of cases go unreported otherwise this number could be much higher (22). Summer season is reported more suitable for this disease when compared to winter (23).

Food can be contaminated by *Campylobacter* in many ways. Mostly the infected animals like cows, turkeys and chickens which do not show any signs and symptoms but serves as carriers and source of pathogen spread (24). *Campylobacter* can reside in several organs of these animals and when this contaminated meat is consumed may lead to infection in human (25). Around 400 million cases of campylobacteriosis have been reported each year worldwide and still there are lot other cases which are not documented especially from developing countries (26). Out of these reported cases 15% have to be hospitalized and 5% of those hospitalized lead to death (27). The consumption of contaminated meat, milk, egg and other by products is considered as main source of infection of *Campylobacter* and other foodborne pathogens (28-34). One can acquire infection by direct contact, contaminated equipment or water (31, 33). The major route of transmission to human is consumption of contaminated food. The animal origin food can be contaminated during slaughtering, dressing or equipment contamination. The other source of infection in modern era is consumption of contaminated ready to eat food products. These food products are mostly contaminated by the infected food handlers or cross contamination during preparation, packaging and storage (35).

The mechanism of disease caused by *Campylobacter* has been investigated worldwide and several mechanism of its pathogenesis has been reported (20, 22). The bacterial flagellum which make it motile and helps to adhere the intestinal mucosa and later invasion into the epithelial cell play a vital role in onset of disease (20).

Among reported 26 species of *Campylobacter*, *Campylobacter jejuni* (*C. jejuni*) is major specie which leads to gastroenteritis in human and is followed by *Campylobacter coli* (*C. coli*) (20). It has been reported that very low dose (360 CFU) of *C. jejuni* can lead to campylobacteriosis and 800 CFU was required to cause diarrhoea in volunteers (36).

Several other studies have reported that onset of disease is dependent on *C. jejuni* strain involved, like strain 81-176 may lead to campylobacteriosis with significantly lower dose when compared with CG8421 strain (36-38).

The epidemiological data is inconclusive as the developing countries of Asia, Africa and Middle East have not adopted the standard reporting protocols (21). However, based on available data it has been observed that the cases of campylobacteriosis have been increased during recent decades (23). Based on available reports, the data suggests the huge variation of cases among different countries and cities of the same country (23). One of the proposed reasons for such variation is the diagnostic method as *Campylobacter* is not that easy to grow so many countries are not equipped with those modern sophisticated tools and techniques to perform the proper diagnosis. The other reason regarding variation in reported cases among different countries was the immune level of population (3). The people with strong immunity can tolerate the infection without showing any sign and symptoms of the disease (13). Importantly, children of developing countries are the most affected population and the occurrence of *campylobacteriosis* decreases significantly with increase in age (38). The possible reasoning is that exposure in early age may lead to protection from reoccurrence of disease. However, these asymptomatic adult carriers might lead to infect the children by direct or indirect contact (38).

The outbreaks of *Campylobacter* have been reported frequently worldwide and the most common cause reported was consumption of contaminated poultry meat and water. Out of 143 outbreaks, 114 were those caused by consumption of poultry meat (21). Similarly, more than 70% of cases of campylobacteriosis in Switzerland were associated with consumption of chicken (25). Similarly in Pakistan meat from different sources was found to be contaminated with *Campylobacter* (39-41) and several other studies have also documented that domesticated animals are also one of the key reservoirs of *Campylobacter* (42-44). The meat consumed from these animals lead to development of disease (44). Moreover, direct contact with these domesticated animals has great potential to infect the ones who come in contact. One study from Denmark reported that 17% of the campylobacteriosis cases were those who were in direct contact with cattle and in Switzerland this figure reached to 19.3% (43). However, there is great variation regarding the prevalence of *Campylobacter* in cattle. It varies from 23 to 90% (44). The most isolated species of *Campylobacter* from cattle are *jejuni*, *coli*, *larai* and *lanienae*. Pigs and wild animals have also been reported as potential reservoirs of *Campylobacter* and can cause disease when their meat is consumed or by direct or indirect contact (45, 46).

Water has also been documented as the source to transmit *Campylobacter* (47). Importantly, both animals and human can be infected after consumption of contaminated water (48). Several studies performed around the globe have reported prevalence of *Campylobacter* in drinking water (49). Furthermore, researchers reported that the people and animals who consumed that contaminated water were found positive for *Campylobacter* (50). So far the water as potential source of *Campylobacter* transmission is underestimated and extensive research is required to explore it in detail.

Developing countries are the most affected ones in term of *Campylobacter* infection in general and especially the food borne (3, 39). The data reported from these countries is nowhere near to real situation as most of the cases go undiagnosed and even several diagnosed cases are not documented and reported properly (3). However the limited studies conducted in these under developed countries have reported that *C. jejuni* is on top of the list as a causative agent of diarrhea (18). The one study conducted at different cities of Bangladesh (Mirzapur), Pakistan (Karachi), India (Kolkata) and African countries found that *C. jejuni* was one of the most common agents in diarrheal cases (18).

In developing countries the most common technique used to detect *Campylobacter* is use of selective media to grow it (18). This technique has several draw backs and the most important one is that growth conditions vary for different *Campylobacter* spp. (18). Thus adopting optimum growth conditions for specific specie will hinder the growth of others. The other modern molecular techniques are employed in developed countries however most of the developing nations cannot provide such facility to all population. The alternate techniques such as enzyme immune assay and PCR based diagnosis have demonstrated promising results but its long journey ahead before adopted by the under developed countries (50). Such poor diagnostic facilities have lead to irrational use of antibiotics and this resulted in increase of *Campylobacter* strains which are resistant to antibiotics (51). Macrolide and fluoroquinolones resistant strains have been increased significantly during the recent decade (51). To cope with such situation European countries banned the use of fluoroquinolones in poultry, however the use of same drug in poultry feed in developing countries is not regulated and unfortunately it is used commonly (52-54).

The control of *Campylobacter* exposure and spread in developing countries is the most difficult task as the contamination of food items and water is very common (2). Furthermore, the unawareness of public has made it almost impossible. To control the spread, it's of utmost important that strict hygienic measure should be implemented at all levels. The breeding and rearing of poultry and animals should be monitored regularly and the birds and animals should be screened for infections before slaughtering (55). The vendors and food handlers should also be screened for infections and certification process may be implemented for those involved in food industry. To provide immunization against *Campylobacter* several attempts have been made to develop a vaccine but so far no concrete success has been achieved (56).

CONCLUSION

In conclusion, the epidemiology and burden of *Campylobacter* is not fully documented, especially in developing countries. The modern molecular techniques should be adopted for the accurate and efficient diagnosis of infection. To minimize the spread, strict hygienic measures must be adopted and the use of antibiotics should be rational.

References:

1. Bintsis T. Foodborne pathogens. *AIMS microbiology*. 2017;3(3):529.
2. Bintsis T. Microbial pollution and food safety. *AIMS microbiology*. 2018;4(3):377.
3. Kirk MD, Pires SM, Black RE, Caipo M, Crump JA, Devleeschauwer B, Döpfer D, Fazil A, Fischer-Walker CL, Hald T, Hall AJ. World Health Organization estimates of the global and regional disease burden of 22 foodborne bacterial, protozoal, and viral diseases, 2010: a data synthesis. *PLoS medicine*. 2015 Dec 3;12(12):e1001921.
4. Thomas MK, Murray R, Flockhart L, Pintar K, Pollari F, Fazil A, Nesbitt A, Marshall B. Estimates of the burden of foodborne illness in Canada for 30 specified pathogens and unspecified agents, circa 2006. *Foodborne pathogens and disease*. 2013 Jul 1;10(7):639-48.
5. Flores GE, Bates ST, Caporaso JG, Lauber CL, Leff JW, Knight R, Fierer N. Diversity, distribution and sources of bacteria in residential kitchens. *Environmental microbiology*. 2013 Feb;15(2):588-96.
6. Cardinale M, Kaiser D, Lueders T, Schnell S, Egert M. Microbiome analysis and confocal microscopy of used kitchen sponges reveal massive colonization by *Acinetobacter*, *Moraxella* and *Chryseobacterium* species. *Scientific reports*. 2017 Jul 19;7(1):1-3.
7. Powell DA, Jacob CJ, Chapman BJ. Enhancing food safety culture to reduce rates of foodborne illness. *Food Control*. 2011 Jun 1;22(6):817-22.
8. Medeiros CO, Cavalli SB, Salay E, Proença RP. Assessment of the methodological strategies adopted by food safety training programmes for food service workers: A systematic review. *Food Control*. 2011 Aug 1;22(8):1136-44.
9. Mensah LD, Julien D. Implementation of food safety management systems in the UK. *Food control*. 2011 Aug 1;22(8):1216-25.
10. Kafetzopoulos DP, Psomas EL, Kafetzopoulos PD. Measuring the effectiveness of the HACCP food safety management system. *Food control*. 2013 Oct 1;33(2):505-13.
11. Dora M, Kumar M, Van Goubergen D, Molnar A, Gellynck X. Food quality management system: Reviewing assessment strategies and a feasibility study for European food small and medium-sized enterprises. *Food control*. 2013 Jun 1;31(2):607-16.
12. Choudhury M, Mahanta L, Goswami J, Mazumder M, Pegoo B. Socio-economic profile and food safety knowledge and practice of street food vendors in the city of Guwahati, Assam, India. *Food Control*. 2011 Feb 1;22(2):196-203.
13. Samapundo S, Climat R, Xhaferi R, Devlieghere F. Food safety knowledge, attitudes and practices of street food vendors and consumers in Port-au-Prince, Haiti. *Food Control*. 2015 Apr 1;50:457-66.
14. Choudhury M, Mahanta LB, Goswami JS, Mazumder MD. Will capacity building training interventions given to street food vendors give us safer food?: A cross-sectional study from India. *Food Control*. 2011 Aug 1;22(8):1233-9.
15. Samapundo S, Thanh TC, Xhaferi R, Devlieghere F. Food safety knowledge, attitudes and practices of street food vendors and consumers in Ho Chi Minh city, Vietnam. *Food Control*. 2016 Dec 1;70:79-89.
16. Aluko OO, Ojeremi TT, Olaleke DA, Ajidagba EB. Evaluation of food safety and sanitary practices among food vendors at car parks in Ile Ife, southwestern Nigeria. *Food Control*. 2014 Jun 1;40:165-71.
17. Asiegbu CV, Lebelo SL, Tabit FT. The food safety knowledge and microbial hazards awareness of consumers of ready-to-eat street-vended food. *Food Control*. 2016 Feb 1;60:422-9.
18. Platts-Mills JA, Kosek M. Update on the burden of *Campylobacter* in developing countries. *Current opinion in infectious diseases*. 2014 Oct;27(5):444.
19. Devleeschauwer B, Bouwknecht M, Manges MJ, Havelaar AH. Health and economic burden of *Campylobacter*. In *Campylobacter* 2017 Jan 1 (pp. 27-40). Academic Press.
20. Moore JE, Corcoran D, Dooley JS, Fanning S, Lucey B, Matsuda M, McDowell DA, Mégraud F, Millar BC, O'Mahony R, O'Riordan L. *Campylobacter*. *Veterinary research*. 2005 May 1;36(3):351-82.
21. Kaakoush NO, Castaño-Rodríguez N, Mitchell HM, Man SM. Global epidemiology of *Campylobacter* infection. *Clinical microbiology reviews*. 2015 Jul 1;28(3):687-720.
22. Epps SV, Harvey RB, Hume ME, Phillips TD, Anderson RC, Nisbet DJ. Foodborne *Campylobacter*: infections, metabolism, pathogenesis and reservoirs. *International journal of environmental research and public health*. 2013 Dec;10(12):6292-304.
23. Hearnden* M, Skelly C, Eyles R, Weinstein P. The regionality of campylobacteriosis seasonality in New Zealand. *International Journal of Environmental Health Research*. 2003 Dec 1;13(4):337-48.

24. Humphrey T, O'Brien S, Madsen M. Campylobacters as zoonotic pathogens: a food production perspective. *International journal of food microbiology*. 2007 Jul 15;117(3):237-57.
25. Eng SK, Pusparajah P, Ab Mutalib NS, Ser HL, Chan KG, Lee LH. Salmonella: a review on pathogenesis, epidemiology and antibiotic resistance. *Frontiers in Life Science*. 2015 Jul 3;8(3):284-93.
26. Garcia S, Heredia NL. 11 Campylobacter. *Guide to foodborne pathogens*. 2013 Jul 10:188.
27. Ruiz-Palacios GM. The health burden of Campylobacter infection and the impact of antimicrobial resistance: playing chicken.
28. Heredia N, García S. Animals as sources of food-borne pathogens: A review. *Animal nutrition*. 2018 Sep 1;4(3):250-5.
29. Samad A, Abbas F, Ahmad Z, Pokryshko O, Asmat TM. Prevalence of foodborne pathogens in food items in Quetta, Pakistan. *Pakistan J. Zool*. 2018 Aug 1;50(4):1-4.
30. Samad A, Abbas F, Ahmad Z, Tanveer Z, Ahmad I, Patching SG, Nawaz N, Asmat MT, Raziq A, Naeem M, Akhtar MA. Multiplex polymerase chain reaction detection of Shiga toxin genes and antibiotic sensitivity of *Escherichia coli* O157: H7 isolated from beef meat in Quetta, Pakistan. *Journal of Food Safety*. 2018 Dec;38(6):e12540.
31. Tareen AM, Ullah K, Asmat TM, Samad A, Iqbal A, Mustafa MZ, Ahmad I. Incidence of Diarrheagenic *Escherichia coli* Pathotypes in Children Suffering from Diarrhea in Tertiary Care Hospitals, Quetta, Pakistan. *Pakistan Journal of Zoology*. 2019 Dec 1;51(6).
32. Samad A, Abbas F, Tanveer Z, Ahmad Z, Asmat TA, Raziq AB, Zahid M. Prevalence of *Salmonella* spp. in chicken meat from Quetta retail outlets and typing through multiplex PCR. *Rom. Biotechnol. Lett.*. 2018;10:1-1.
33. Mengal H, Samad A, Asmat MT, Mustafa MZ, Abbas F, Sajjad N, Ishtiaq H, Rahim H, Ghilzai D. 3. The study to evaluate the prevalence of *Yersinia* species in salad sold and water supplied in Quetta city, Pakistan. *Pure and Applied Biology (PAB)*. 2019 Aug 15;8(3):1902-8.
34. Asmat TM. Molecular detection of *Escherichia coli* (*E. coli*) from Diarrheal stool samples from children in Quetta, Balochistan, Pakistan. *Pak-Euro Journal of Medical and Life Sciences*. 2019 Dec 14;2(2):27-31.
35. Igwaran A, Okoh AI. Human campylobacteriosis: a public health concern of global importance. *Heliyon*. 2019 Nov 1;5(11):e02814.
36. Nachamkin I. Chronic effects of Campylobacter infection. *Microbes and infection*. 2002 Apr 1;4(4):399-403.
37. Hendrixson DR, DiRita VJ. Identification of Campylobacter jejuni genes involved in commensal colonization of the chick gastrointestinal tract. *Molecular microbiology*. 2004 Apr;52(2):471-84.
38. Thielman NM, Guerrant RL. Acute infectious diarrhea. *New England Journal of Medicine*. 2004 Jan 1;350(1):38-47.
39. Hussain I, Mahmood MS, Akhtar M, Khan A. Prevalence of Campylobacter species in meat, milk and other food commodities in Pakistan. *Food microbiology*. 2007 May 1;24(3):219-22.
40. Nisar M, Mushtaq MH, Shehzad W, Hussain A, Nasar M, Nagaraja KV, Goyal SM. Occurrence of Campylobacter in retail meat in Lahore, Pakistan. *Acta tropica*. 2018 Sep 1;185:42-5.
41. Khalil K, Lindblom GB, Mazhar K, Sjögren E, Kaijser B. Frequency and enterotoxigenicity of Campylobacter jejuni and *C. coli* in domestic animals in Pakistan as compared to Sweden. *The Journal of tropical medicine and hygiene*. 1993 Feb;96(1):35-40.
42. Sheppard SK, Cheng L, Méric G, De Haan CP, Llerena AK, Marttinen P, Vidal A, Ridley A, Clifton-Hadley F, Connor TR, Strachan NJ. Cryptic ecology among host generalist Campylobacter jejuni in domestic animals. *Molecular ecology*. 2014 May;23(10):2442-51.
43. Sheppard SK, Maiden MC. The evolution of Campylobacter jejuni and Campylobacter coli. *Cold Spring Harbor perspectives in biology*. 2015 Aug 1;7(8):a018119.
44. Manyi-Loh CE, Mamphweli SN, Meyer EL, Makaka G, Simon M, Okoh AI. An overview of the control of bacterial pathogens in cattle manure. *International journal of environmental research and public health*. 2016 Sep;13(9):843.
45. Rahimi E, Ameri M, Kazemeini HR. Prevalence and antimicrobial resistance of Campylobacter species isolated from raw camel, beef, lamb, and goat meat in Iran. *Foodborne pathogens and disease*. 2010 Apr 1;7(4):443-7.
46. Uaboi-Egbenni PO, Bessong PO, Samie S, Obi CL. Prevalence and antimicrobial susceptibility profiles of Campylobacter jejuni and coli isolated from diarrheic and non-diarrheic goat faeces in Venda region, South Africa. *African Journal of Biotechnology*. 2011;10(64):14116-24.

47. Miller WG, Mandrell RE. Prevalence of *Campylobacter* in the food and water supply: incidence, outbreaks, isolation and detection. *Campylobacter: Molecular and cellular biology*. 2005 Jun 1:101-63.
48. Wilkes G, Edge TA, Gannon VP, Jokinen C, Lyautey E, Neumann NF, Ruecker N, Scott A, Sunohara M, Topp E, Lapen DR. Associations among pathogenic bacteria, parasites, and environmental and land use factors in multiple mixed-use watersheds. *Water Research*. 2011 Nov 15;45(18):5807-25.
49. Nachamkin I. *Campylobacter jejuni*. In *Food Microbiology: Fundamentals and Frontiers*, Third Edition 2007 Jan 1 (pp. 237-248). American Society of Microbiology.
50. Vidic J, Manzano M, Chang CM, Jaffrezic-Renault N. Advanced biosensors for detection of pathogens related to livestock and poultry. *Veterinary research*. 2017 Dec;48(1):1-22.
51. Doyle MP, Diez-Gonzalez F, Hill C, editors. *Food microbiology: fundamentals and frontiers*. John Wiley & Sons; 2020 Jul 10.
52. Economou V, Gousia P. Agriculture and food animals as a source of antimicrobial-resistant bacteria. *Infection and drug resistance*. 2015;8:49.
53. Bolton DJ. *Campylobacter* virulence and survival factors. *Food microbiology*. 2015 Jun 1;48:99-108.
54. Chen X, Naren GW, Wu CM, Wang Y, Dai L, Xia LN, Luo PJ, Zhang Q, Shen JZ. Prevalence and antimicrobial resistance of *Campylobacter* isolates in broilers from China. *Veterinary microbiology*. 2010 Jul 29;144(1-2):133-9.
55. Umaraw P, Prajapati A, Verma AK, Pathak V, Singh VP. Control of campylobacter in poultry industry from farm to poultry processing unit: A review. *Critical reviews in food science and nutrition*. 2017 Mar 4;57(4):659-65.
56. Dhama K, Rajagunalan S, Chakraborty S, Verma AK, Kumar A, Tiwari R, Kapoor S. Food-borne pathogens of animal origin-diagnosis, prevention, control and their zoonotic significance: a review. *Pakistan journal of biological sciences*. 2013 Oct 15;16(20):1076.